




Scan me!



Mining waste - source of environmental pollution

Akmaral Kuldaeva¹  [orcid: 0009-0003-1377-171X](https://orcid.org/0009-0003-1377-171X)
e-mail: akmaral.kuldaeva@mail.ru

Musabek Duvanakov²  [orcid: 0009-0009-4068-7584](https://orcid.org/0009-0009-4068-7584)
e-mail: mduvanakov@mail.ru

^{1,2}Osh technological university named after M.M.Adyshev, Osh. 714018, Republic of Kyrgyzstan.

<https://doi.org/10.5281/zenodo.15176517>

Abstract: In this article mining wastes and their negative impact are considered. The main wastes and tailing ponds of the mining industry located on the territory of Kyrgyzstan are the wastes of mining enterprises since the times of the Soviet Union. At present, the problems of these tailings are of regional and global nature. The aim of the study is to give a geo-ecological assessment of this problem.

Key words: waste, tailings, mining, mining industry, geoecology, pollution of nature, uranium, ore, etc.

1 Introduction

Most of the territory of Kyrgyzstan is located at an altitude of more than 2500 m above sea level. The territory below 1500 m is only 1/8 of the country's territory. Permafrost covers 4% of the country's area (8,000 km²) and stores 650 billion cubic metres of fresh water. Also, the subsoil of the mountains of Kyrgyzstan is rich in various mineral resources, and their exploitation makes a significant contribution to the economic development of the country. For example, gold production alone accounts for 11 per cent of the country's GDP, 40 per cent of industry and 48 per cent of exports.

However, in accordance with the goal of sustainable development, it is necessary to take into account not only the economic side of the problem, but also the environmental side.

As we know, it is necessary to emphasise that all mining operations are harmful industrial facilities. In particular, mining wastes containing various toxic and radioactive substances are very dangerous for the environment. In general, as a result of all activities related to mining (exploration - extraction - processing, metallurgical processing, etc.), a large amount of rocks and slags generated during production are excavated and accumulated on the surface of the earth.

2 Relevance of the problems and research materials

In addition, the current stage of development of the mining industry is due to the fact that the amount of mineral-rich ores is decreasing and the technology of extraction of valuable minerals is gradually improving (more precisely, valuable minerals are extracted as environmentally friendly). and if possible with the help of new technologies). This process increases the number of storage facilities for waste treatment.

One of the most dangerous environmental problems of Kyrgyzstan, as well as other Central Asian countries, began with the mining and production of uranium and toxic chemical elements (antimony, silicon, vanadium and nickel).

Today Kyrgyzstan is one of the most environmentally dangerous regions of the planet in terms of radioactive poisoning. In a relatively small territory of the country, there are 49 uranium waste storage sites and 80 accumulation sites of processed rock. In these places 70 million m³ of highly toxic radioactive waste is buried, which corresponds to 14 m³ for each inhabitant of the republic [1]. Depending on the type and concentration of the mined ores, these waste repositories contain various fine radionuclides, heavy metal compounds harmful to health, and toxic substances used as reagents in mineral extraction. The latter include cyanides, acids, sulphates, nitrates and others.

Currently, the largest waste storage facility in the country is the Kumtor cyanide waste storage facility with a total design volume of 100 million m³ ha. The fact that these Kumtor waste storage facilities are located in the headwaters of the Naryn River and in the glacier zone creates a very dangerous situation.

Waste metals such as antimony and mercury are disposed of in sludge pits and slag ponds. However, the amount of waste from sorting antimony and mercury is relatively small compared to other mining operations. For example, the volume of accumulated waste at the Aidarken and Kadamjai plants is 15 million m³. They also occupy many territories and damage the environment to a certain extent.

Large amounts of such anthropogenic wastes on the earth surface are located in some regions of the country (Mailuu-Suu, Sumsar, Kan, Kadamzhai, Ak-Tuz, etc.) have created an extremely dangerous environmental situation at the regional level [2].

About groundwater contamination in the area of Kara-Balta mine tailings dump location (Kara-Balta) The analyses carried out showed that the area of contaminated water extended up to 15 km and the contamination was very intense and persistent. It was also known that groundwater contamination had moved towards the city and low-lying settlements [3]. Alarming, groundwater and groundwater contamination has led to a reduction in uranium mining and continues unabated, even as the amount of waste in waste storage facilities decreases.

A critical geo-ecological situation has developed in the regions where most mining enterprises operate. Here the most acute problem is the pollution of nature and deterioration of human health in large settlements [2].

Forced storage of a huge amount of waste, its removal, transportation, neutralisation, storage, burial requires huge financial costs, alienates territories for their disposal, and disturbs the ecology as a whole [4].

The majority of waste storage facilities in Kyrgyzstan are located in the basins of the Naryn, Mailuu-Suu and Chui rivers, so they are transboundary in nature. Out of 36 waste storage facilities, 31 contain radioactive waste, including 28 direct uranium production wastes (total volume 8.2 million m³), 3 thorium-containing polymetal production wastes. Non-ferrous metal and chemical production waste in 5 landfills (5.15 million m³). Waste from the former lead-zinc plant is stored in 3 waste repositories in Sumsar. Here, according to the Ministry of Health and Social Welfare, the amount of cadmium exceeds the norm 320 times and manganese 9 times [5].

The total exposure power of each uranium waste repository is 30-100 thousand $\mu\text{R/s}$. And in the upper part of waste repositories it ranges from 27 to 60 $\mu\text{R/s}$, in some places this indicator reaches 600-1000 $\mu\text{R/s}$, exceeding the MPC by 20-50 times. Thus, in general, artificial gamma radiation in these zones is possible. Thus, the annual radiation rate is 5 times more per person [6].

Long-term protection measures against various natural phenomena (landslides, heavy rainfall, floods and inundations) were not provided during the construction of these waste storage facilities. In addition, no sanitary protection zones were organised to protect settlements. As a result, the territories of the republics of Kyrgyzstan, Kazakhstan, Tajikistan and Uzbekistan, where more than 5 million people live, remain in acute environmental danger [7].

3 Conclusion

In general, the following factors determine the emergence of the crisis situation in the region:

- mining extracts rocks from the lower layer of the earth's crust to the surface, and toxic zinc, lead, manganese, sometimes considered particularly toxic;

- mercury, arsenic, cadmium, and easily soluble or completely insoluble minerals are also extracted from it. As these rocks undergo the process of destruction, the volume or surface area increases by a factor of 3-10 and accumulates as waste at the earth's surface, and the filtration coefficient increases by a factor of 100-120;
- as mentioned above, the surface area and filtration coefficient increased several times, the waste rock poor in composition (valuable components are separated) becomes more convenient for precipitation and runoff. It is easily washed away and drains away together with ground and surface waters;
- as a result of the above processes there is intensive pollution of ground and surface waters;
- in addition to polluting the hydrosphere, surface residues are spread by the wind and pollute the environment with dust. Thus, toxic compounds in the form of dust harm the soil and plants.

The most dangerous part of mining waste accumulating on the surface is that its harmful effects last for a very long time. For example, according to geochemical monitoring at Canada's Elliot Lake mine, mine tailings have been contaminating groundwater for 120 years. No one knows how long these contaminations will last.

According to calculations by the World Bank, the TACIS project and Kyrgyz experts, the country will need \$24 million to solve the problems with radioactive waste storage facilities. Firstly, it is difficult for the current budget of the country to allocate such an amount at once. Secondly, we believe that solving these issues, which have a regional scale, concerns not only Kyrgyzstan. This is due to the fact that almost all of the current waste storage sites are remnants of mines mined during the former Soviet Union. Also, the mined mineral resources were used in the interests of the countries that were part of the Soviet Union. Therefore, this should be a problem of the immediate neighbourhood.

Reference

1. Bykovchenko Y.G. et al. Technogenic uranium pollution of the biosphere of Kyrgyzstan // Monograph. - Bishkek, 2005. -171 c.
2. Torgoev I.A. Geo-ecological monitoring at development of resources of mountains of Kyrgyzstan. -Bishkek: Exponenta, 2000.
3. National Environmental Protection Plan. Priorities 1995 - 1997. -Bishkek, 1995. -82 c.
4. Duvanakulov M.A. Modern methods of analysis of ecological problems on development of non-metallic materials // M.A. Duvanakulov/ Science, new technologies and innovations of Kyrgyzstan № 9, 2023. -Б., 2023. - 35-38 c.
5. Geo-ecological safety and risk of natural-technogenic catastrophes on the territory of Kyrgyzstan. Torgoev I.A., Aleshin Y.G., Moldobaeva B.B. -Bishkek: Zheka Ltd, 1999.
6. Berdibaeva A.B. Impact of uranium tailings on biosphere objects in Kyrgyzstan // Bykovchenko Y.G., Zhunushov A.T., Tukhvatshin R.R. / Medicine in Kyrgyzstan № 2. -Б., 2014. -132-133 c.
- 7.https://unece.org/fileadmin/DAM/env/documents/2019/TEIA/Workshop_Nov_2019/Session_5_RUS/1a_Ms._Gulfiya_Shabayeva_KYR_needs_and_challenges_RUS.pdf